

Bicyclic graphs with exactly two main signless Laplacian eigenvalues *

HE HUANG, HANYUAN DENG[†]

College of Mathematics and Computer Science,

Hunan Normal University, Changsha, Hunan 410081, P. R. China

(October 30, 2011)

Abstract

A signless Laplacian eigenvalue of a graph G is called a main signless Laplacian eigenvalue if it has an eigenvector the sum of whose entries is not equal to zero. In this paper, all connected bicyclic graphs with exactly two main eigenvalues are determined.

Keywords: Spectra of a graph, main signless Laplacian eigenvalue, bicyclic graph.

AMS classification: 05C50

1 Introduction

Let G be a simple connected graph with vertex set $V = \{v_1, v_2, \dots, v_n\}$ and $(0, 1)$ -adjacency matrix A . An eigenvalue λ of A is said to be a main eigenvalue of G if the eigenspace $\varepsilon(\lambda)$ is not orthogonal to the all-1 vector $\mathbf{j}$, i.e., it has an eigenvector the sum of whose entries is not equal to zero. An eigenvector $\mathbf{x}$ is a main eigenvector if $\mathbf{x}^T \mathbf{j} \neq 0$. For details of

*This work was supported by Hunan Provincial Natural Science Foundation of China (09JJ6009) and the Program for Science and Technology Innovative Research Team in Higher Educational Institution of Hunan Province.

[†]Corresponding author: hydeng@hunnu.edu.cn.

the main eigenvector of adjacency matrix, the readers are suggested to refer to [1] [3] [4] [5] [6] [7].

Very recently, we introduced in [8] the main signless Laplacian eigenvalue of a graph. Let $L^+ = D + A$, where $D = \text{diag}(d(v_1), d(v_2), \dots, d(v_n))$ is the diagonal matrix of vertex degrees and $d(v_i)$ is the degree of vertex v_i . Then L^+ is called the signless Laplacian matrix of G . Some results on the signless Laplacian matrix of a graph can be found in [9] [10] [11] [12] [13]. An eigenvalue μ of L^+ is said to be a main signless Laplacian eigenvalue of G (or a main eigenvalue of L^+) if the eigenspace $\varepsilon(\mu)$ is not orthogonal to the all-1 vector $\mathbf{j}$, i.e., it has an eigenvector the sum of whose entries is not equal to zero. It was showed in [8] that a graph G with exactly one main signless Laplacian eigenvalue if and only if G is regular, and the trees, unicyclic graphs with exactly two main signless Laplacian eigenvalues were characterized, respectively.

A graph G is called a bicyclic graph, if it is a simple connected graph in which the number of edges equals the number of vertices plus one. The aim of this work is to characterize all bicyclic graphs with exactly two main signless Laplacian eigenvalues.

2 Some Lemmas

In [8], an alternative characterization of a graph with exactly two main signless Laplacian eigenvalues is given by 2-walks.

A graph G is called 2-walk (a, b) -**parabolic** [8] if there exist uniquely a positive integer a and a non-negative integer b such that $a^2 - 8b > 0$ and $s(v) = -d^2(v) + ad(v) - b$ holds for all $v \in V(G)$, where $s(v) =$

$\sum_{u \in N_G(v)} d(u)$ and $N_G(v)$ is the set of all neighbors of v in G .

Note that any 2-walk (a, b) -parabolic graph is irregular, since one has $s(v) = -d^2(v) + 2kd(v) - 0$ and $s(v) = -d^2(v) + (2k + 1)d(v) - k$ for a k -regular graph, i.e., $(a, b) = (2k, 0)$ or $(2k + 1, k)$ is not unique.

Lemma 1 [8]. A graph G has exactly two main signless Laplacian eigenvalues if and only if there exist uniquely a positive integer a and a non-negative integer b such that $a^2 - 8b > 0$ and

$$s(v) = -d^2(v) + ad(v) - b \quad (1)$$

for all $v \in V(G)$, i.e., G is a 2-walk (a, b) -parabolic graph.

In [8], we showed that a graph with exactly one main signless Laplacian eigenvalue if and only if it is regular. So, a 2-walk (a, b) -parabolic graph G is non-regular, and there are $u, v \in V(G)$ such that $d(u) \neq d(v)$. From $s(u) = -d^2(u) + ad(u) - b$ and $s(v) = -d^2(v) + ad(v) - b$, we have

$$\begin{aligned} a &= \frac{s(u) - s(v)}{d(u) - d(v)} + d(u) + d(v) \\ b &= \frac{s(u) - s(v)}{d(u) - d(v)} d(v) + d(u)d(v) - s(v) \end{aligned} \quad (2)$$

Remark 1. If G is a 2-walk (a, b) -parabolic graph with $\delta(G) = 1$, then $a - b \geq 3$ since there is a pendent vertex x with the only incident edge xy in G and $d(y) = s(x) = -1 + a - b \geq 2$.

Let $\mathcal{G}_{a,b} = \{G : G \text{ is a 2-walk } (a, b)\text{-parabolic bicyclic graph with } \delta(G) = 1\}$. For each $G \in \mathcal{G}_{a,b}$, G_0 is the graph obtained from G by deleting all pendent vertices. If $v \in V(G_0)$, we use $d_{G_0}(v)$ to denote the degree of the vertex v in G_0 . The following lemmas are true for unicyclic graphs, and it is easy to see that they are also true for bicyclic graphs from their proofs in [8].

Lemma 2 [8]. If $G \in \mathcal{G}_{a,b}$ and $v \in V(G_0)$, then $d(v) = d_{G_0}(v)$ or $d(v) = a - b - 1$.

Lemma 3 [8]. If $G \in \mathcal{G}_{a,b}$, then (i) $\delta(G_0) \geq 2$; (ii) $a - b \geq 4$ and $a \geq 5$.

Lemma 4. Let $G \in \mathcal{G}_{a,b}$ and $C = u_1 u_2 \dots u_k u_1$ a cycle of G . If $d_{G_0}(u_1) \geq 3$ and $d_{G_0}(u_2) = 2$, then there is $i \in \{1, 2, \dots, k\}$ such that $d(u_i) \neq a - b - 1$.

Proof. By the definition of G_0 , $C = u_1 u_2 \dots u_k u_1$ is also a cycle of G_0 .

If $d(u_1) = d(u_2) = \dots = d(u_k) = a - b - 1$, then from (1)

$$s(u_1) = -(a - b - 1)^2 + a(a - b - 1) - b = s(u_2) \quad (3)$$

And

$$\begin{aligned} s(u_1) &= d(u_1) - d_{G_0}(u_1) + d(u_2) + d(u_k) + \sum_{w \in N_{G_0}(u_1) \setminus \{u_2, u_k\}} d(w) \\ &\geq 3(a - b - 1) + 2(d_{G_0}(u_1) - 2) - d_{G_0}(u_1) \\ &= 3(a - b - 1) + d_{G_0}(u_1) - 4 \\ &\geq 3(a - b - 1) - 1 \quad (\text{by } d_{G_0}(u_1) \geq 3) \\ s(u_2) &= d(u_1) + d(u_3) + d(u_2) - d_{G_0}(u_2) \quad (\text{by } d_{G_0}(u_2) = 2) \\ &= 3(a - b - 1) - 2 \end{aligned}$$

So, $s(u_1) > s(u_2)$, it contradicts (3). $\square$

If R is a cycle or a path of G , then the length of R , denoted by $l(R)$, is defined as the number of edges of R .

Lemma 5. Let G be a 2-walk (a, b) -parabolic graph. $R = x_1 x_2 \dots x_t$ is a path or cycle of length at least 2 in G such that $d(x_1) \geq 3$, $d(x_t) \geq 3$ and $d(x_2) = \dots = d(x_{t-1}) = 2$. Then

(i) If $d(x_1) = d(x_t)$, then $l(R) \leq 3$; Moreover, if $l(R) = 3$, then there exists no path $Q = y_1 y_2 y_3$ in G such that $d(y_1) = d(y_3) = d(x_1)$ and

$$d(y_2) = 2;$$

(ii) If $d(x_1) \neq d(x_t)$, then $l(R) \leq 2$.

Proof. (i) If $l(R) \geq 4$, then $d(x_2) = d(x_3) = d(x_4) = 2$. By (1), we have $s(x_2) = d(x_3) + d(x_1) = -2^2 + 2a - b = s(x_3) = d(x_2) + d(x_4) = 4$, and $d(x_1) = 2$. This contradicts $d(x_1) \geq 3$. So, $l(R) \leq 3$.

If $l(R) = 3$, and there is a path $Q = y_1 y_2 y_3$ in G such that $d(y_1) = d(y_3) = d(x_1)$ and $d(y_2) = 2$, then by (1), $s(y_2) = d(y_1) + d(y_3) = -2^2 + 2a - b = s(x_2) = d(x_1) + d(x_3) = d(x_1) + 2$, and $d(y_1) + d(y_3) = d(x_1) + 2$. This is impossible since $d(y_1) = d(y_3) = d(x_1) \geq 3$.

(ii) If $l(R) \geq 3$, then $s(x_2) = -2^2 + 2a - b = s(x_{t-1})$ by (1). On the other hand, $s(x_2) = d(x_1) + d(x_3) = d(x_1) + 2$ and $s(x_{t-1}) = d(x_{t-2}) + d(x_t) = d(x_t) + 2$, a contradiction with $d(x_1) \neq d(x_t)$. $\square$

Lemma 6. Let $G \in \mathcal{G}_{a,b}$, $R = u_1 u_2 \cdots u_k$ a path or cycle of G_0 with length at least 3 such that $d_{G_0}(u_1) = d_{G_0}(u_k) \in \{3, 4\}$ and $d_{G_0}(u_i) = 2$ for $2 \leq i \leq k-1$.

(i) If there is $i \in \{2, \dots, k-1\}$ such that $d(u_i) \neq d(u_{i+1})$, then $l(R) = 3$ and $b = 1$. Moreover,

If $d_{G_0}(u_1) = 3$, then $a = 6$, $d(u_1) = d_{G_0}(u_1) = 3$, $d(u_2) = 2$, $d(u_3) = 4$; or $d(u_2) = 4$, $d(u_3) = 2$.

If $d_{G_0}(u_1) = 4$, then $a = 7$, $d(u_1) = d_{G_0}(u_1) = 4$, $d(u_2) = 2$, $d(u_3) = 5$; or $d(u_2) = 5$, $d(u_3) = 2$.

(ii) If $d(u_2) = d(u_3) = \cdots = d(u_{k-1})$, then $d(u_2) = d(u_3) = \cdots = d(u_{k-1}) \in \{2, a - b - 1\}$. Moreover, if R is a cycle, then $l(R) = 3$ and $d(u_2) = d(u_3) = 2$.

Proof. (i) By Lemma 2, $d(u_j) \in \{d_{G_0}(u_j), a - b - 1\} = \{2, a - b - 1\}$,

$2 \leq j \leq k-1$. If there is $i \in \{2, 3, \dots, k-2\}$ such that $d(u_i) \neq d(u_{i+1})$, then $d(u_i) = 2$ and $d(u_{i+1}) = a-b-1$, or $d(u_{i+1}) = 2$ and $d(u_i) = a-b-1$. Without loss of generality, assume that $d(u_i) = 2$, $d(u_{i+1}) = a-b-1$. From (2), we have

$$\begin{aligned} a &= \frac{s(u_{i+1}) - s(x)}{d(u_{i+1}) - d(x)} + d(u_{i+1}) + d(x) \\ &= \frac{2 + d(u_{i+2}) + a - b - 3 - (a - b - 1)}{a - b - 1 - 1} + a - b - 1 + 1 \end{aligned}$$

where x is a pendent vertex of G , and

$$d(u_{i+2}) = b(a-b-2) \tag{4}$$

$b \neq 0$ since $d(u_{i+2}) \neq 0$. So, $b \geq 1$.

Case 1. $b = 1$. Then $d(u_{i+2}) = a-3 < a-2 = a-b-1$, and $d(u_{i+2}) = d_{G_0}(u_{i+2})$ from Lemma 2. If $u_{i+2} \neq u_k$, then $d(u_{i+2}) = d_{G_0}(u_{i+2}) = 2$. So, $a = 5$, $a-b-1 = 3$. Because $d(u_1) \geq d_{G_0}(u_1) \in \{3, 4\}$ and $d(u_1) \in \{d_{G_0}(u_1), a-b-1\} = \{d_{G_0}(u_1), 3\}$, by Lemma 2, we have $d(u_1) = d_{G_0}(u_1)$ and $N_G(u_1) = N_{G_0}(u_1)$.

From (1), $s(u_1) = -d_{G_0}^2(u_1) + 5d_{G_0}(u_1) - 1$. On the other hand, $s(u_1) = \sum_{w \in N_G(u_1)} d(w) = \sum_{w \in N_{G_0}(u_1)} d(w) \geq 2d_{G_0}(u_1)$. So,

$$-d_{G_0}^2(u_1) + 5d_{G_0}(u_1) - 1 \geq 2d_{G_0}(u_1), \text{ i.e., } d_{G_0}^2(u_1) - 3d_{G_0}(u_1) + 1 \leq 0$$

This implies $\frac{3-\sqrt{5}}{2} \leq d_{G_0}(u_1) \leq \frac{3+\sqrt{5}}{2} < 3$, a contradiction with $d_{G_0}(u_1) \geq 3$.

So, $u_{i+2} = u_k$.

Since $d(u_i) = 2$, $s(u_i) = -2^2 + 2a - b = 2a - 5$ from (1), and $d(u_{i-1}) = s(u_i) - d(u_{i+1}) = 2a - 5 - (a - 2) = a - 3 \neq a - b - 1$. By Lemma 2, $d_{G_0}(u_{i-1}) = d(u_{i-1}) = a - 3 = d(u_{i+2}) = d(u_k) > 2$. So, $u_{i-1} = u_1$, and $l(R) = 3$. It implies $i = 2$.

If $d_{G_0}(u_1) = 3$, then $d_{G_0}(u_1) = d_{G_0}(u_{i-1}) = d(u_{i-1}) = a - 3$ from above, and $a = 6$. $d(u_2) = d(u_i) = 2$ and $d(u_3) = d(u_{i+1}) = a - b - 1 = 4$.

Similarly, if $d_{G_0}(u_1) = 4$, then $a - 3 = 4$, and $a = 7$. $d(u_2) = 2$ and $d(u_3) = 5$.

Case 2. $b = 2$.

If $d_{G_0}(u_k) = 3$, then from (4) and Lemma 3, $d(u_{i+2}) = 2(a - b - 2) = a - b - 1 + a - b - 3 > a - b - 1 \geq 3 \geq d_{G_0}(u_{i+2})$. This contradicts Lemma 2 whether $R = u_1u_2 \cdots u_k$ a path or cycle.

If $d_{G_0}(u_k) = 4$, then from (4) and Lemma 3, $d(u_{i+2}) = 2(a - b - 2) > a - b - 1 > 2$. By Lemma 2, we have $d_{G_0}(u_{i+2}) = d(u_{i+2}) > 2$. So, $u_{i+2} = u_k$, and $d(u_k) = d_{G_0}(u_k) = 4$. This implies $2(a - b - 2) = 4$ and $a = 6$.

From (1), $s(u_k) = -d^2(u_k) + ad(u_k) - b = 6$. On the other hand, $s(u_k) = \sum_{w \in N_G(u_k)} d(w) = \sum_{w \in N_{G_0}(u_k)} d(w) \geq 2d_{G_0}(u_k) = 8$, a contradiction.

Case 3. $b > 2$.

From (4), $d(u_{i+2}) = b(a - b - 2) \geq 3(a - b - 2) = a - b - 1 + 2(a - b - 2) - 1 > \max\{a - b - 1, d_{G_0}(u_{i+2})\}$. This contradicts Lemma 2.

(ii) It is obvious that $d(u_2) = \cdots = d(u_{k-1}) \in \{2, a - b - 1\}$ from Lemma 2, since $d(u_2) = \cdots = d(u_{k-1})$ and $d_{G_0}(u_2) = \cdots = d_{G_0}(u_{k-1}) = 2$.

Moreover, if R is a cycle and $d(u_2) = \cdots = d(u_{k-1}) = a - b - 1$, then $d(u_1) \neq a - b - 1$ from Lemma 4, and $d(u_1) = d_{G_0}(u_1)$ from Lemma 2. By (1), $s(u_2) = -(a - b - 1)^2 + a(a - b - 1) - b$. On the other hand,

$s(u_2) = d(u_1) + d(u_3) + d(u_2) - d_{G_0}(u_2) = 2(a - b - 1) - 2 + d(u_1)$. So,

$$\begin{aligned}
d(u_1) &= s(u_2) - 2(a - b - 1) + 2 \\
&= -(a - b - 1)^2 + a(a - b - 1) - b - 2(a - b - 1) + 2 \\
&= (b - 1)(a - b - 1) - b + 2 \\
&= (b - 1)(a - b - 2) + 1
\end{aligned}$$

If $b \leq 1$, then $d(u_1) \leq 1$. This is impossible.

If $b = 2$, then $d(u_1) = a - b - 1$. This contradicts $d(u_1) \neq a - b - 1$.

If $b > 2$, then $d(u_1) \geq 2(a - b - 2) + 1 \geq 5 > d_{G_0}(u_1)$. But $d(u_1) \neq a - b - 1$ from Lemma 4, it contradicts Lemma 2.

So, $d(u_2) = \dots = d(u_{k-1}) = 2$. And $l(R) = 3$ from Lemma 5(i). $\square$

Lemma 7. Let $G \in \mathcal{G}_{a,b}$, $u, v \in V(G_0)$ and $d(u) = d(v)$. If $N_{G_0}(u) = \{u', u''\}$ and $N_{G_0}(v) = \{v', v''\}$, then $d(u') = d(v')$ if and only if $d(u'') = d(v'')$.

Proof. From (1) and $d(u) = d(v)$, we have

$$s(u) = -d^2(u) + ad(u) - b = -d^2(v) + ad(v) - b = s(v)$$

On the other hand,

$$\begin{aligned}
s(u) &= d(u') + d(u'') + d(u) - d_{G_0}(u) \\
s(v) &= d(v') + d(v'') + d(v) - d_{G_0}(v)
\end{aligned}$$

So, $d(u') = d(v')$ if and only if $d(u'') = d(v'')$. $\square$

3 Bicyclic graphs with exactly two main signless Laplacian eigenvalues

In the following, we determine all bicyclic graphs with exactly two main signless Laplacian eigenvalues.

Theorem 1. There is no bicyclic graph with exactly two main signless Laplacian eigenvalues and minimum degree one, i.e., $\mathcal{G}_{a,b} = \emptyset$.

Proof. By way of contradiction, assume that $G \in \mathcal{G}_{a,b}$. From Lemma 3, $\delta(G_0) \geq 2$. So, G_0 is one of the three graphs F_1, F_2, F_3 shown in Figure 1.

Case 1. $G_0 = F_1$, where $|V(C_1) \cap V(C_2)| = 1$, $C_1 = u_1 u_2 \cdots u_n$, $C_2 = v_1 v_2 \cdots v_m$ and $u_1 = u_n = v_1 = v_m = u$.

From Lemma 6, we know that $n = m = 4$ and (i) if $d(u_2) = d(u_3)$, $d(v_2) = d(v_3)$, then $G = F_{11}$; (ii) if $d(u_2) = d(u_3)$, $d(v_2) \neq d(v_3)$ (or $d(u_2) \neq d(u_3)$, $d(v_2) = d(v_3)$), then $G = F_{12}$; (iii) if $d(u_2) \neq d(u_3)$, $d(v_2) \neq d(v_3)$, then $G = F_{13}$.

Note that $G \in \mathcal{G}_{a,b}$ and $\delta(G) = 1$, there is a pendant vertex x in G .

If $G = F_{11}$, then $d(u_2) = d(u_3) = d(v_2) = d(v_3) = 2$, $d(u_1) = a - b - 1$. By (2),

$$a = \frac{s(u_2) - s(x)}{d(u_2) - d(x)} + d(u_2) + d(x) = \frac{a - b - 1 + 2 - (a - b - 1)}{2 - 1} + 2 + 1 = 5$$

and

$$\begin{aligned} a &= \frac{s(u_1) - s(x)}{d(u_1) - d(x)} + d(u_1) + d(x) = \frac{8 + a - b - 1 - 4 - (a - b - 1)}{a - b - 1 - 1} + a - b - 1 + 1 \\ &= \frac{4}{a - b - 2} + a - b \end{aligned}$$

So, $b^2 - 3b + 4 = 0$. This is impossible.

If $G = F_{12}$ or F_{13} , then $a = 7$ and $b = 1$ by Lemma 6(i). Using Eq.(1), it is easy to check that $G = F_{12}$ and F_{13} are not 2-walk (a, b) -parabolic graphs, a contradiction.

Case 2. $G_0 = F_2$, where $V(C_1) \cap V(C_2) = \emptyset$, $C_1 = u_1 u_2 \cdots u_n$, $C_2 = v_1 v_2 \cdots v_m$, $P = w_1 w_2 \cdots w_t$, $u_1 = u_n = w_1 = u$ and $w_t = v_1 = v_m = v$.

From Lemma 6, we know that $n = m = 4$ and (i) if $d(u_2) \neq d(u_3)$, $d(v_2) = d(v_3)$ (or $d(u_2) = d(u_3)$, $d(v_2) \neq d(v_3)$), then $G \cong F_{21}$; (ii) if $d(u_2) \neq d(u_3)$, $d(v_2) \neq d(v_3)$, then $G \cong F_{22}$; (iii) if $d(u_2) = d(u_3)$, $d(v_2) = d(v_3)$, then $G \cong F_{23}$.

(i) In the graph F_{21} , we have $a = 6$, $b = 1$, $d(u_2) = 2$, $d(u_3) = 4$, $d(u_1) = 3$, $d(v_2) = 2$, $d(v_3) = 2$ by Lemma 6. $d(v_1) = 5$ by using Eq.(1) to v_2 , and it can be found that a pendant vertex x of v_1 does not satisfy Eq.(1). So, F_{21} is not a 2-walk (a, b) -parabolic graph.

(ii) In the graph F_{22} , we have $a = 6$, $b = 1$, $d(u_2) = 2$, $d(u_3) = 4$, $d(u_1) = 3$, $d(v_2) = 2$, $d(v_3) = 4$, $d(v_1) = 3$ by Lemma 6. First, $d(w_2) = 2$ by using Eq.(1) to u_1 , and then $d(w_3) = 4$ by using Eq.(1) to w_2 , and finally $d(w_4) = 3$ by using Eq.(1) to w_3 . If $w_4 = v_1$, then $s(v_1) = d(w_3) + d(v_2) + d(v_3) = 4 + 2 + 4 \neq -d^2(v_1) + ad(v_1) - b$, i.e., v_1 does not satisfy Eq.(1); otherwise, a pendant vertex x of w_4 does not satisfy Eq.(1). So, F_{22} is not a 2-walk (a, b) -parabolic graph.

(iii) In the graph F_{23} , $d(u_2) = d(u_3) = d(v_2) = d(v_3) = 2$.

First, we show that $d(u_1) = d(v_1) = 3$. If $d(u_1) \neq 3$, then $d(u_1) > d_{G_0}(u_1) = 3$ and $d(u_1) = a - b - 1$ from Lemma 2. By Eq.(1), we have $s(u_2) = -2^2 + 2a - b = 2a - b - 4$. On the other hand, $s(u_2) = d(u_1) + d(u_3) = a - b - 1 + 2$. So, $a - b - 1 + 2 = 2a - b - 4$ and $a = 5$. It implies that $b = 0$ since b is a nonnegative integer and $a - b - 1 = d(u_1) > d_{G_0}(u_1) = 3$, and $d(u_1) = a - b - 1 = 4$. Using Eq.(1) again, $s(u_1) = -d^2(u_1) + ad(u_1) - b = -16 + 20 - 0 = 4 < d(u_2) + d(u_3) + d(w_2) + (d(u_1) - d_{G_0}(u_1))$, a contradiction. So, $d(u_1) \neq 3$. Similarly (or by symmetry), $d(v_1) = 3$.

Now, $s(u_2) = -2^2 + 2a - b = 2a - 4 - b$ by Eq.(1), and $s(u_2) = d(u_1) +$

$d(u_3) = 5$. So, $2a - b = 9$. And $a = 5$, $b = 1$ from Lemma 3. Using Eq.(1) again, we have $s(u_1) = 5 = d(u_2) + d(u_3) + d(w_2)$, and $d(w_2) = 1$, a contradiction.

So, F_{23} is not a 2-walk (a, b) -parabolic graph.

Case 3. $G_0 = F_3$, where $P_1 = u_1u_2 \cdots u_n$, $P_2 = v_1v_2 \cdots v_m$, $P_3 = w_1w_2 \cdots w_t$, $u_1 = v_1 = w_1 = u$, $u_n = v_m = w_t = v$. And we may assume that $n, m \geq 3$ since G is a simple graph.

Claim 1. $d(u_2) = \cdots = d(u_{n-1})$, $d(v_2) = \cdots = d(v_{m-1})$, $d(w_2) = \cdots = d(w_{t-1})$.

If there $i \in \{2, \cdots, n-2\}$ such that $d(u_i) \neq d(u_{i+1})$, then $l(P_1) = 3$ (i.e., $n = 4$), $b = 1$, $a = 6$, $d(u_1) = d_{G_0}(u_1) = 3$, $d(u_4) = d_{G_0}(u_4) = 3$, $d(u_2) = 2$, $d(u_3) = 4$ or $d(u_2) = 4$, $d(u_3) = 2$. By symmetry, we may assume that $d(u_2) = 2$, $d(u_3) = 4$. Using Eq.(1), $s(u_4) = 8 = d(u_3) + d(v_{m-1}) + d(w_{t-1})$ implies $d(v_{m-1}) = d(w_{t-1}) = 2$. It can be obtained that $d(v_{m-2}) = d(w_{t-2}) = 4$ by using Lemma 7 for $\{u_2, v_{m-1}\}$ and $\{u_2, w_{t-1}\}$, respectively. And $l(P_2) = l(P_3) = 3$ (i.e., $m = t = 4$) from Lemma 6(i). Now, $s(u_1) = d(u_2) + d(v_2) + d(w_2) = 10 \neq -d^2(u_1) + ad(u_1) - b$, it contradicts Eq.(1). So, $d(u_2) = \cdots = d(u_{n-1})$.

Similarly, $d(v_2) = \cdots = d(v_{m-1})$ and $d(w_2) = \cdots = d(w_{t-1})$. $\square$

Claim 2. $d(u_1) = d(u_n)$.

By way of contradiction, assume that $d(u_1) \neq d(u_n)$. $d(u_1), d(u_n) \in \{3, a - b - 1\}$ from Lemma 2 and $d_{G_0}(u_1) = d_{G_0}(u_n) = 3$. By symmetry, we may assume that $d(u_1) = a - b - 1$, $d(u_n) = 3$. Then $a - b - 1 \geq 4$, i.e.,

$$a - b \geq 5 \tag{5}$$

From Claim 1, $d(u_2) = d(u_{n-1})$, $d(v_2) = d(v_{m-1})$, $d(w_2) = d(w_{t-1})$. And

$$s(u_1) - s(u_n) = \begin{cases} \{(d(u_1) - d_{G_0}(u_1)) + d(u_2) + d(v_2) + d(u_n)\} \\ - \{d(u_{n-1}) + d(v_{m-1}) + d(u_1)\}, & l(P_3) = 1; \\ \{(d(u_1) - d_{G_0}(u_1)) + d(u_2) + d(v_2) + d(w_2)\} \\ - \{d(u_{n-1}) + d(v_{m-1}) + d(w_{t-1})\}, & l(P_3) \geq 2 \end{cases}$$

$$= \begin{cases} 0, & l(P_3) = 1; \\ a - b - 4, & l(P_3) \geq 2. \end{cases}$$

From Eq.(2), we have

$$a = \frac{s(u_1) - s(u_n)}{d(u_1) - d(u_n)} + d(u_1) + d(u_n) = \begin{cases} a - b + 2, & l(P_3) = 1; \\ a - b + 3, & l(P_3) \geq 2. \end{cases}$$

So,

$$b = \begin{cases} 2, & l(P_3) = 1; \\ 3, & l(P_3) \geq 2. \end{cases}$$

From Eq.(2) again,

$$s(u_1) = -(a - b - 1)^2 + a(a - b - 1) - b = (b + 1)(a - b - 1) - b.$$

On the other hand, $s(u_1) = (d(u_1) - d_{G_0}(u_1)) + d(u_2) + d(v_2) + d(w_2)$.

So,

$$\begin{aligned} d(u_2) + d(v_2) + d(w_2) &= s(u_1) - d(u_1) + d_{G_0}(u_1) = b(a - b - 1) - b + 3 \\ &= \begin{cases} 2(a - b - 1) + 1, & l(P_3) = 1; \\ 3(a - b - 1), & l(P_3) \geq 2. \end{cases} \end{aligned} \quad (6)$$

By Lemma 2,

$$\begin{aligned} d(u_2) &\in \{d_{G_0}(u_2), a - b - 1\} = \{2, a - b - 1\} \\ d(v_2) &\in \{d_{G_0}(v_2), a - b - 1\} = \{2, a - b - 1\} \\ d(w_2) &\in \{d_{G_0}(w_2), a - b - 1\} = \{2, 3, a - b - 1\} \end{aligned}$$

If $l(P_3) \geq 2$, then Eq.(6) implies $d(u_2) = d(v_2) = d(w_2) = a - b - 1$;

If $l(P_3) = 1$, then $w_2 = u_n$ and $d(w_2) = 3$. Eq.(6) implies $d(u_2) + d(v_2) = (a - b - 1) + (a - b - 3)$. So, $\{d(u_2), d(v_2)\} = \{2, a - b - 1\}$. Without loss of generality, we assume that $d(u_2) = a - b - 1$.

In the following, we show that $l(P_1) = 2$. Otherwise, $a - b - 1 = d(u_2) = d(u_3) = \dots = d(u_{n-1})$ from Claim 1. By Eq.(1), $s(u_2) = s(u_{n-1})$ since $d(u_2) = d(u_{n-1})$. But $s(u_2) = (d(u_2) - d_{G_0}(u_2)) + d(u_3) + d(u_1)$ and $s(u_{n-1}) = (d(u_2) - d_{G_0}(u_2)) + d(u_{n-2}) + d(u_n)$, it implies $d(u_1) = d(u_n)$, a contradiction. So, $l(P_1) = 2$, and $n = 3$, $d(u_3) = d(u_n) = 3$.

Now, by (1), $s(u_2) = -(a - b - 1)^2 + a(a - b - 1) - b = (b + 1)(a - b - 1) - b$.

On the other hand,

$$\begin{aligned} s(u_2) &= d(u_1) + d(u_3) + d(u_2) - d_{G_0}(u_2) \\ &= (a - b - 1) + 3 + (a - b - 1 - 2) = 2(a - b - 1) + 1. \end{aligned}$$

So, $(b + 1)(a - b - 1) - b = 2(a - b - 1) + 1$, and $(b - 1)(a - b - 2) = 2$. This is impossible since $b \geq 2$ and $a - b \geq 5$ from Eq.(5). $\square$

Claim 3. If $d(u_1) = d(u_n) = a - b - 1$, then at most one of u_2, v_2, w_2 has degree $a - b - 1$.

Otherwise, we may assume that $d(u_2) = d(v_2) = a - b - 1$. By Claim 1, $d(v) = a - b - 1$ for any $v \in V(P_1) \cup V(P_2) \setminus \{u_1, u_n\}$. Then, all vertices on the cycle $C = P_1 \cup P_2 = u_1 u_2 \dots u_n v_{m-1} \dots v_2 u_1$ have degree $a - b - 1$, a contradiction to Lemma 4. $\square$

Claim 4. $d(u_1) = d(u_n) = 3$.

We only need to prove that $d(u_1) = 3$ by Claim 2. If $d(u_1) \neq 3 = d_{G_0}(u_1)$,

then $d(u_1) = a - b - 1$ by Lemma 2, and

$$a - b - 1 > 3, \text{ i.e., } a - b \geq 5. \quad (7)$$

From Claim 3, at least two of u_2, v_2, w_2 have degree 2. Without loss of generality, assume that $d(u_2) = d(v_2) = 2$. By Eq.(1), $s(u_1) = -(a - b - 1)^2 + a(a - b - 1) - b = (b + 1)(a - b - 1) - b$. On the other hand, $s(u_1) = (d(u_1) - d_{G_0}(u_1)) + d(u_2) + d(v_2) + d(w_2) = (a - b) + d(w_2)$ and $2 \leq d(w_2) \leq a - b - 1$. We have

$$(a - b - 1) + 3 \leq (b + 1)(a - b - 1) - b \leq 2(a - b - 1) + 1$$

and (i) $b + 3 \leq b(a - b - 1)$, implying that $b \neq 0$ and $b \geq 1$; (ii) $(b - 1)(a - b - 2) \leq 2$, implying that $b \leq 1$. So, $b = 1$.

By Eq.(1), $s(u_2) = -2^2 + 2a - b = 2a - b - 4$. And $d(u_3) = s(u_2) - d(u_1) = a - 3 < a - b - 1$, $d(u_3) = d_{G_0}(u_3) = 2$ from Lemma 2 and $u_3 \neq u_n$ since $d(u_n) = d(u_1) = a - b - 1$. So, $a = d(u_3) + 3 = 5$. But now $a - b = 4$ contradicts Eq.(7). Thus, $d(u_1) = 3$. $\square$

In the following, we will show that G is not a 2-walk (a, b) -parabolic graph if $G_0 = F_3$ by Claims 1-4.

If there is a 2-walk (a, b) -parabolic graph G such that $G_0 = F_3$, let x be a pendent vertex of G and xy is the only edge incident with x . Then, $y \in (V(P_1) \cup V(P_2) \cup V(P_3)) \setminus \{u_1, u_n\}$ by Claim 4 and Lemma 3, $d_{G_0}(y) = 2$ and $d(y) = a - b - 1$. Let $N_{G_0}(y) = \{y', y''\}$, $4 \leq d_{G_0}(y') + d_{G_0}(y'') \leq d(y') + d(y'') \leq 2(a - b - 1)$ by Lemma 2. By Eq.(1), $s(y) = d(y') + d(y'') + (d(y) - d_{G_0}(y)) = -(a - b - 1)^2 + a(a - b - 1) - b = (b + 1)(a - b - 1) - b$, i.e., $d(y') + d(y'') = b(a - b - 2) + 2$. So,

$$4 \leq b(a - b - 2) + 2 \leq 2(a - b - 1).$$

The left inequality implies $b > 0$, and the right inequality implies that $(b - 2)(a - b - 2) \leq 0$, i.e., $b \leq 2$, since $a - b - 2 > 0$ by Lemma 3. So, $b = 1$ or $b = 2$.

Without loss of generality, we assume $y \in V(P_1) \setminus \{u_1, u_n\}$. By Claim 1, $d(u_2) = \cdots = d(u_{n-1}) = d(y) = a - b - 1$. Using Eq.(1),

$$\begin{aligned} a &= \frac{s(u_2) - s(x)}{d(u_2) - d(x)} + d(u_2) + d(x) \\ &= \frac{(d(u_1) + d(u_3) + d(u_2) - d_{G_0}(u_2)) - (a - b - 1)}{a - b - 1 - 1} + (a - b - 1) + 1 \\ &= \frac{d(u_3) + 1}{a - b - 2} + a - b \end{aligned}$$

and

$$d(u_3) = b(a - b - 2) - 1 \quad (8)$$

(i) If $b = 1$, then by Eq.(8), $d(u_3) = a - 4 < a - b - 1$, and $u_3 = u_n$, $d(u_3) = d(u_n) = 3$. So, $a = 7$. By Eq.(1), we have $s(u_1) = d(u_2) + d(v_2) + d(w_2) = -d^2(u_1) + ad(u_1) - b = -9 + 21 - 1 = 11$, and $d(v_2) + d(w_2) = s(u_1) - d(u_2) = 11 - 5 = 6$. Since $d(v_2), d(w_2) \in \{d_{G_0}(v_2), d_{G_0}(w_2), a - b - 1\}$ by Lemma 2 and $a - b - 1 = 5$, $d(v_2) = d_{G_0}(v_2) \leq 3$ and $d(w_2) = d_{G_0}(w_2) \leq 3$. So, $d(v_2) = d_{G_0}(v_2) = 3$, $d(w_2) = d_{G_0}(w_2) = 3$, and $v_2 = u_n = w_2$, i.e., $m = t = 2$, contradicts that G is a simple graph.

(ii) If $b = 2$, then by Eq.(8), $d(u_3) = 2(a - b - 2) - 1 = (a - b - 1) + (a - b - 3) > a - b - 1 \geq d_{G_0}(u_3)$, contradicts Lemma 2.

So, there is not a 2-walk (a, b) -parabolic graph G such that $G_0 = F_3$, i.e., G is not a 2-walk (a, b) -parabolic graph if $G_0 = F_3$.

The proof of Theorem 1 is completed. $\square$

Theorem 1 also shows that a bicyclic graph G with the minimum degree $\delta(G) = 1$ has at least three main signless Laplacian eigenvalues.

The following result characterizes bicyclic graphs with exactly two main signless Laplacian eigenvalues.

Theorem 2. A bicyclic graph G has exactly two main signless Laplacian eigenvalues, i.e., G is a 2-walk (a, b) -parabolic graph if and only if G is isomorphic to one of $G_1, G_2, \dots, G_7$ (see Figure 4).

Proof. Let G be a bicyclic graph. If G is a 2-walk (a, b) -parabolic graph, then $\delta(G) > 1$ from Theorem 1. So, G is one of the three graphs F_1, F_2, F_3 shown in Figure 1.

Case 1. $G = F_1$. Then $l(C_1) = l(C_2) = 3$ from Lemma 6(ii), i.e., $G = G_1$. Using Eq.(1), it is easy to check that G_1 is a 2-walk $(7, 4)$ -parabolic graph.

Case 2. $G = F_2$. Then $l(C_1) = l(C_2) = 3$ from Lemma 6(ii), and $l(P) = 1$ or $l(P) = 3$ from Lemma 5(i). So, $G = G_2$ or $G = G_3$. It can be checked easily that G_2 and G_3 are 2-walk $(7, 5)$ -parabolic and $(6, 3)$ -parabolic graphs, respectively.

Case 3. $G = F_3$. Without loss of generality, we assume that $l(P_1) \geq l(P_2) \geq l(P_3)$. Then $l(P_1) \geq l(P_2) \geq 2$ since G is a simple graph. By Lemma 5(i), $l(P_i) \leq 3$ ($i = 1, 2, 3$) and if one of $l(P_1), l(P_2), l(P_3)$ is 3, then the other two are not 2. So, $G = G_4$ or G_5 if $l(P_3) = 1$; $G = G_6$ if $l(P_3) = 2$; $G = G_7$ if $l(P_3) = 3$. Also, G_4, G_5, G_6, G_7 are 2-walk $(6, 2)$ -parabolic, $(7, 5)$ -parabolic, $(5, 0)$ -parabolic and $(6, 3)$ -parabolic graphs.

Therefore, a bicyclic graph G with exactly two main signless Laplacian eigenvalues if and only if G is isomorphic to one of $G_1, G_2, \dots, G_7$.




Figure 1. The graphs F_1, F_2, F_3 .




Figure 2. The graphs F_{11}, F_{12}, F_{13} , where $r = 0$ or $r = a - b - 5$.




Figure 3. The graphs F_{21}, F_{22}, F_{23} .




Figure 4.

All bicyclic graphs with exactly two main signless Laplacian eigenvalues.

References

- [1] D. Cvetković, P. Rowlinson, S. Simić, Eigenspaces of Graphs, Cambridge University Press, Cambridge, 1997.
- [2] D. Cvetković, M. Peterić, A table of connected graphs on six vertices, Discrete Math. 50 (1984) 37-49.
- [3] E. M. Hagos, Some results on graph spectra, Linear Algebra Appl. 356 (2002) 103-111.
- [4] Y. Hou, H. Zhou, Trees with exactly two main eigenvalues, Acta of Hunan Normal University, 28 (2) (2005) 1-3 (in Chinese).
- [5] Y. Hou, F. Tian, Unicyclic graphs with exactly two main eigenvalues, Appl. Math. Lett. 19 (2006) 1143-1147.
- [6] Z. Hu, S. Li, C. Zhu, Bicyclic graphs with exactly two main eigenvalues, Linear Algebra Appl. 431 (2009) 1848-1857.
- [7] P. Rowlinson, The main eigenvalues of a graph: a survey, Applicable Analysis and Discrete Mathematics, 1 (2007) 445-471.
- [8] H. Deng, H. Huang, On the main signless Laplacian eigenvalues of a graph, submitted.
- [9] D. Cvetković, Signless Laplacians and line graphs, Bull. Acad. Serbe Sci. Arts, Cl. Sci. Math. Natur., Sci. Math. 131(30) (2005) 85-92.
- [10] D. Cvetković, P. Rowlinson, S. K. Simić, Signless Laplacians of finite graphs, Linear Algebra Appl., 423(2007) 155-171.

- [11] D. Cvetković, P. Rowlinson, S. K. Simić, Eigenvalue bounds for the signless Laplacian, *Publ. Inst. Math. (Beograd)*, 81(95)(2007) 11-27.
- [12] D. Cardoso, D. Cvetković, P. Rowlinson, S. K. Simić, A sharp lower bound for the least eigenvalue of the signless Laplacian of a non-bipartite graph, *Linear Algebra Appl.*, 429(2008) 2770-2780.
- [13] D. Cvetković, New theorems for signless Laplacian eigenvalues, *Bull. Acad. Serbe Sci. Arts, Cl. Sci. Math. Natur., Sci. Math.* 137(33) (2008) 131-146.
- [14] K. Feng, *Algebraic Number Theory*, Beijing, Science Press, 2000 (in Chinese).